A•Cute Kids Urgent Care

Tylenol (acetaminophen) Dosage Chart

Give the weight appropriate dosage listed below every 4 hours as needed for a temperature > 101 °F.

*****Do not give Tylenol to children under 3 months of age without first consulting a doctor.**

Weight (pounds)	Infant Drops 80mg / 0.8ml	Childrens Liquid 160mg / 5ml	Chewable Tablets 80mg tablets	Junior Strength 160 mg tablets	For Staff Use: Weight (Kg)
6 -7	40 mg (0.4 ml)				3
8-9	60 mg (0.6 ml)				4
10-11	75 mg (0.8 ml)				5
12-13	90 mg (0.9 ml)				6
14-16	100 mg (1.0 ml)				7
17-18	120 mg (1.2 ml)				8
19-22	135 mg (1.4 ml)				9
22-26	160 mg (1.6 ml)	5 ml (1 tsp)	2 tabs	1 tab/cap	10-12
28-31		6 ml	2 tabs	1 tab/cap	13-14
32-35		7 ml (1 ½ tsp)	3 tabs	1 ½ tabs/caps	15-16
36-39		8 ml	3 tabs	1 ½ tabs/caps	17-18
40-44		9 ml	3 tabs	1 ½ tabs/caps	19-20
45-48		10 ml (2 tsp)	4 tabs	2 tabs/caps	21-22
49-53		11 ml	4 tabs	2 tabs/caps	23-24
54-57		12 ml (2 ½ tsp)	5 tabs	2 ½ tabs/caps	25-26
58-62		13 ml	5 tabs	2 ½ tabs/caps	27-28
63-66		14 ml	5 tabs	2 ½ tabs/caps	29-30
67-71		15 ml (3 tsp)	6 tabs	3 tabs/caps	31-32
72-75		16 ml	6 tabs	3 tabs/caps	33-34
76-80		17 ml (3 ½ tsp)	7 tabs	3 ½ tabs/caps	35-36
81-84		18 ml	7 tabs	3 ½ tabs/caps	37-38
85-88		19 ml	7 tabs	3 ½ tabs/caps	39-40
89-93		20 ml (4 tsp)	8 tabs	4 tabs/caps	41-42
94-99		20 ml (4 tsp)	8 tabs	4 tabs/caps	43-44
≥ 100		20 ml (4 tsp)	8 tabs	4 tabs/caps	≥ 45

When used together, tylenol and motrin may be alternated every 3 hours.

****Consult a physician before using tylenol if your child has any liver problems.***

A•Cute Kids Urgent Care

Ibuprofen (Motrin, Advil) Dosage Chart

Give the weight appropriate dosage listed below every 6 hours as needed for temperature > 101 °F or for pain.

*****Do not give Ibuprofen if child is less than 6 months old*****

Weight (pounds)	Infant Drops 50mg / 1.25 ml	Childrens Liquid 100mg / 5ml	For Staff Use Weight (Kg)
8-9	40 mg (1.0 ml)		4
10-11	50 mg (1.25 ml)		5
12-13	60 mg (1.5 ml)		6
14-16	70 mg (1.75ml)		7
17-18	80 mg (2.0 ml)		8
19-22	90 mg (2.25ml)		9
22-26	100 mg (2.5ml)	5 ml (1 tsp)	10-11
28-31	120 mg (3.0ml)	6 ml	12-13
32-35	140 mg (3.5ml)	7 ml (1 ½ tsp)	14-15
36-39	160 mg (4.0 ml)	8 ml	16-17
40-44		9 ml	18-19
45-48		10 ml (2 tsp)	20-21
49-53		11 ml	22-23
54-57		12 ml (2 ½ tsp)	24-25
58-62		13 ml	26-27
63-66		14 ml	28-29
67-71		15 ml (3 tsp)	30-31
72-75		16 ml	32-33
76-80		17 ml (3 ½ tsp)	34-35
81-84		18 ml	36-37
85-88		19 ml	38-39
89-93		20 ml (4 tsp)	40-41
94-99		20 ml (4 tsp)	42-43
≥ 100		20 ml (4 tsp)	≥ 44

When used together, tylenol and motrin may be alternated every 3 hours.

****Consult a physician before using motrin if your child has any kidney problems.***